

Quasi-Projective Reduction of Toric Varieties

Annette A'Campo-Neuen and Jürgen Hausen

Abstract

We define a quasi-projective reduction of a complex algebraic variety X to be a regular map from X to a quasi-projective variety that is universal with respect to regular maps from X to quasi-projective varieties. A toric quasi-projective reduction is the analogous notion in the category of toric varieties. For a given toric variety X we first construct a toric quasi-projective reduction. Then we show that X has a quasi-projective reduction if and only if its toric quasi-projective reduction is surjective. We apply this result to characterize when the action of a subtorus on a quasi-projective toric variety admits a categorical quotient in the category of quasi-projective varieties.

Introduction

Let X be a complex algebraic variety. We call a regular map $p: X \rightarrow X^{\text{qp}}$ from X to a quasi-projective variety X^{qp} a *quasi-projective reduction* of X if every regular map $f: X \rightarrow Z$ to a quasi-projective variety Z factors uniquely through p , i.e., there exists a unique regular map $\tilde{f}: X^{\text{qp}} \rightarrow Z$ such $f = \tilde{f} \circ p$.

As a first result of the present article we characterize when a given toric variety X has a quasi-projective reduction. In Section 1 we construct a *toric quasi-projective reduction* of X , i.e., a toric morphism $q: X \rightarrow X^{\text{tqp}}$ to a quasi-projective toric variety X^{tqp} such that every toric morphism from X to a quasi-projective toric variety factors uniquely through q . Then we prove (see Section 2):

Theorem 1. *A toric variety X has a quasi-projective reduction if and only if its toric quasi-projective reduction $q: X \rightarrow X^{\text{tqp}}$ is surjective. If q is surjective, then it is the quasi-projective reduction of X .*

The above theorem implies in particular that every complete toric variety has a projective reduction. But as we show by an explicit example (see 3.1), the quasi-projective reduction

need not exist in general. We apply Theorem 1 to obtain a complete answer to the following problem, posed by A. Białynicki-Birula:

Let X be a quasi-projective toric variety with acting torus T and let $H \subset T$ be a subtorus. When does the action of H admit a quotient in the category of quasi-projective varieties, i.e., an H -invariant regular map $s: X \rightarrow Y$ to a quasi-projective variety Y such that every H -invariant regular map from X to a quasi-projective variety factors uniquely through s ?

In order to state our answer, let $s_1: X \rightarrow X_{\text{tor}}/H$ denote the toric quotient (see [1]). Recall that s_1 is universal with respect to H -invariant toric morphisms. Moreover, let $q: X_{\text{tor}}/H \rightarrow Y$ be the toric quasi-projective reduction. Then our result is the following (for the proof see Section 2):

Theorem 2. *The action of H on X admits a quotient in the category of quasi-projective varieties if and only if $s := q \circ s_1$ is surjective. If s is surjective, then it is the quotient for the action of H on X .*

Examples of quasi-projective toric varieties with a subtorus action admitting a quotient in the category of quasi-projective varieties are obtained from Mumford's Geometric Invariant Theory. In 3.4 and 3.5 we discuss examples of subtorus actions that have no such quotient.

Notation

A *toric variety* is a normal algebraic variety X endowed with an effective regular action of an algebraic torus T that has an open orbit. We refer to T as the *acting torus* of X . For every toric variety X we fix a base point x_0 in the open orbit.

A regular map $f: X \rightarrow X'$ of toric varieties with base points x_0 and x'_0 respectively is called a *toric morphism* if $f(x_0) = f(x'_0)$ and there is a homomorphism $\varphi: T \rightarrow T'$ of the acting tori such that $f(t \cdot x) = \varphi(t) \cdot f(x)$ holds for every $(t, x) \in T \times X$.

The basic construction in the theory of toric varieties is to associate to a given fan Δ in an n -dimensional lattice an n -dimensional toric variety X_Δ . The assignment $\Delta \mapsto X_\Delta$ is in fact an equivalence of categories (see e.g. [6] or [9]). For our construction of toric quasiprojective reductions we need the following generalization of the notion of a fan:

Let N denote a n -dimensional lattice and set $N_{\mathbb{R}} := \mathbb{R} \otimes_{\mathbb{Z}} N$. A *quasi-fan* in N is a finite set Δ of rational convex polyhedral cones in $N_{\mathbb{R}}$ such that for each $\sigma \in \Delta$ also every face of σ is an element of Δ and any two cones of Δ intersect in a common face. So a quasi-fan is a fan if all its cones are strictly convex. For a quasi-fan Δ , we denote by $\Delta^{\max}$ the set of its maximal cones and by $|\Delta| := \bigcup_{\sigma \in \Delta} \sigma$ its support.

For a homomorphism $F: N \rightarrow N'$ of lattices, let $F_{\mathbb{R}}$ denote the associated homomorphism of real vector spaces. A *map of quasi-fans* Δ in N and Δ' in N' is a lattice homomorphism $F: N \rightarrow N'$ such that for every $\sigma \in \Delta$ there is a $\sigma' \in \Delta'$ with $F_{\mathbb{R}}(\sigma) \subset \sigma'$. As mentioned above, every map $F: \Delta \rightarrow \Delta'$ of fans gives rise to a toric morphism $f: X_{\Delta} \rightarrow X_{\Delta'}$.

Every quasi-fan Δ in N defines in a canonical manner a fan: Let V denote the intersection of all cones of Δ . Then V is a linear subspace of $N_{\mathbb{R}}$. Set $L := V \cap N$ and let $Q: N \rightarrow \tilde{N} := N/L$ denote the projection. Then the cones $Q_{\mathbb{R}}(\sigma)$, $\sigma \in \Delta^{\max}$, are the maximal cones of a fan $\tilde{\Delta}$ in $\tilde{N}$. We call $\tilde{\Delta}$ the *quotient fan* of Δ . By construction, Q is a map of the quasi-fans Δ and $\tilde{\Delta}$.

1 Construction of the Toric Quasi-Projective Reduction

The construction of the toric quasi-projective reduction is done in the category of fans. Toric morphisms from a complete toric variety X_{Δ} to projective spaces are related to concave support functions of the fan Δ . Since we also want to consider non-complete fans it is more natural to work with the following notion instead of support functions:

Let Δ be a quasi-fan in a lattice N . A finite family $\mathfrak{U} := (u_i)_{i \in I}$ of linear forms $u_i \in M := \text{Hom}(N, \mathbb{Z})$ is called Δ -*concave*, if it satisfies the following condition: for every $\sigma \in \Delta^{\max}$ there is an index $i(\sigma)$ such that

$$u_{i(\sigma)}|_{\sigma} \leq u_i|_{\sigma} \quad \text{for all } i \in I.$$

Note that for two given Δ -concave families $\mathfrak{U} := (u_i)_{i \in I}$ and $\mathfrak{U}' := (u'_j)_{j \in J}$ of linear forms the *sum family*

$$\mathfrak{U} + \mathfrak{U}' := (u_i + u'_j)_{(i,j) \in I \times J}$$

is again a Δ -concave family. For a Δ -concave family $\mathfrak{U}$ let $P_{\mathfrak{U}}$ denote the convex hull of $\mathfrak{U}$. Then $P_{\mathfrak{U}}$ is a lattice polytope in $M_{\mathbb{R}}$. Let $\Sigma_{\mathfrak{U}}$ denote the normal quasi-fan of $P_{\mathfrak{U}}$ in N . Recall that the faces P' of P correspond order-reversingly to the cones of $\Sigma_{\mathfrak{U}}$ by

$$P' \mapsto \tau_{P'} := \{v \in N_{\mathbb{R}}; p'(v) \leq p(v) \text{ for all } (p', p) \in P' \times P\}.$$

If $u_1, \dots, u_r$ denote the vertices of $P_{\mathfrak{U}}$, then the family $(u_i)_{i=1, \dots, r}$ is *strictly* $\Sigma_{\mathfrak{U}}$ -concave, i.e., on every relative interior $\tau_{\{u_i\}}^\circ$ the linear form u_i is strictly smaller than the forms u_j with $j \neq i$.

Now assume that Δ is a fan. Call a subset R of the set $\Delta^{(1)}$ of extremal rays of Δ *indecomposable*, if for every Δ -concave family $\mathfrak{U}$ the set R is contained in some maximal cone of $\Sigma_{\mathfrak{U}}$. Let $R_1, \dots, R_k$ be the maximal indecomposable subsets of $\Delta^{(1)}$.

1.1 Lemma. *There exists a Δ -concave family $\mathfrak{U}$ such that every R_i is the intersection of $\Delta^{(1)}$ with some maximal cone of $\Sigma_{\mathfrak{U}}$.*

Proof. For every decomposable subset S of $\Delta^{(1)}$ choose a Δ -concave family $\mathfrak{U}_S$ such that S is not contained in any maximal cone of $\Sigma_{\mathfrak{U}_S}$. Let $\mathfrak{U}$ be the sum of these families $\mathfrak{U}_S$. Then $P_{\mathfrak{U}}$ is the Minkowski-Sum of the $P_{\mathfrak{U}_S}$. Consequently $\Sigma_{\mathfrak{U}}$ is the common refinement of the $\Sigma_{\mathfrak{U}_S}$. This readily yields the claim. $\square$

A Δ -concave family $\mathfrak{U}$ with the property of Lemma 1.1 will be called *generic*. As a consequence of the above lemma we obtain the following statement for the sets

$$\varrho_i := \operatorname{conv} \left(\bigcup_{\varrho \in R_i} \varrho \right).$$

1.2 Remark. The ϱ_i are the maximal cones of a quasi-fan Σ in N . The lattice homomorphism id_N is a map of the quasi-fans Δ and Σ . Moreover, if $\mathfrak{U}$ is a generic Δ -concave family, then id_N is an affine map of the quasi-fans Σ and $\Sigma_{\mathfrak{U}}$. $\square$

Here a map F of quasi-fans Δ in N and Δ' in N' is called *affine* if for every maximal cone σ' of Δ' the set $F_{\mathbb{R}}^{-1}(\sigma') \cap |\Delta|$ is a (maximal) cone of Δ . Note that a map of fans is affine if and only if the associated toric morphism is affine.

Now we construct the quasi-projective toric reduction of a toric variety X_{Δ} defined by the fan Δ . Let V denote the minimal cone of the quasi-fan Σ determined by Δ as in Remark 1.2. Set $L := N \cap V$, let $Q: N \rightarrow \tilde{N} := N/L$ be the projection and denote by $\tilde{\Delta}$ the quotient-fan of Σ .

1.3 Proposition. *The toric morphism $q: X_{\Delta} \rightarrow X_{\tilde{\Delta}}$ associated to Q is the toric quasi-projective reduction of X_{Δ} .*

Proof. First we show that $X_{\tilde{\Delta}}$ is in fact quasi-projective. Choose a generic Δ -concave family $\mathfrak{U} = (u_{\sigma})_{\sigma \in \Delta^{\max}}$. Let V_1 denote the minimal cone of the quasi-fan $\Sigma_{\mathfrak{U}}$. Set $L_1 :=$

$N \cap V_1$, let $P: N \rightarrow \overline{N} := N/L_1$ be the projection and denote the quotient-fan of $\Sigma_{\mathfrak{U}}$ by $\overline{\Delta}$.

Since $\mathfrak{U}$ induces a strictly $\overline{\Delta}$ -concave family, the associated toric variety $X_{\overline{\Delta}}$ is projective. The minimal cone V of Σ is contained in V_1 , so we obtain a lattice homomorphism $G: \tilde{N} \rightarrow \overline{N}$ with $G \circ Q = P$. By construction, G is an affine map of the fans $\tilde{\Delta}$ and $\overline{\Delta}$. So the associated toric morphism $g: X_{\tilde{\Delta}} \rightarrow X_{\overline{\Delta}}$ is affine. Since $X_{\overline{\Delta}}$ is projective we can use [4], Chap. II, Th. 4.5.2, to conclude that $X_{\tilde{\Delta}}$ is quasi-projective.

Now we verify the universal property of q . Let $f: X_{\Delta} \rightarrow X'$ be a toric morphism to a quasi-projective toric variety X' . We may assume that f arises from a map $F: N \rightarrow N'$ of fans Δ and Δ' . Choose a polytopal completion Δ'' of Δ' . By suitable stellar subdivisions (see [5], p. 72) we achieve that every maximal cone of Δ'' contains at most one maximal cone of Δ' .

Let $(u_{\sigma''})_{\sigma'' \in \Delta''^{\max}}$ be a strictly Δ'' -concave family. Then the linear forms $u_{\sigma''} \circ F$ form a Δ -concave family. Let $\sigma \in \Sigma^{\max}$. By construction, σ is mapped by $F_{\mathbb{R}}$ into some cone of Δ'' . Moreover, σ is the convex hull of certain extremal rays of Δ , so $F_{\mathbb{R}}(\sigma)$ is in fact contained in a maximal cone of Δ' . Hence F is a map of the quasi-fans Σ and Δ' . In particular we have $F(L) = 0$. Thus there is a map $\tilde{F}: \tilde{N} \rightarrow N'$ of the fans $\tilde{\Delta}$ and Δ' with $F = \tilde{F} \circ Q$. The associated toric morphism $\tilde{f}: X_{\tilde{\Delta}} \rightarrow X'$ yields the desired factorization of f . $\square$

2 Proof of the Theorems

Let X be a toric variety with acting torus T and assume that $H \subset T$ is an algebraic subgroup. Let Z be an arbitrary quasi-projective variety. We need the following decomposition result for regular maps:

2.1 Proposition. *Let $f: X \rightarrow Z$ be an H -invariant regular map. Then there exist a locally closed subvariety W of some $\mathbb{P}_r$, an H -invariant toric morphism $g: X \rightarrow \mathbb{P}_r$ with $g(X) \subset W$ and a regular map $h: W \rightarrow Z$ such that $f = h \circ g$.*

Proof. In a first step we consider the special case that $Z = \mathbb{P}_m$ and X is an open toric subvariety of some $\mathbb{C}^n$. Then there are polynomials $f_0, \dots, f_m \in \mathbb{C}[z_1, \dots, z_n]$ having no common zero in X such that $f(z) = [f_0(z), \dots, f_m(z)]$ holds for every $z \in X$. Clearly we may assume that the f_i have no non-trivial common divisor.

Since f is H -invariant, every f_i/f_j is an H -invariant rational function on $\mathbb{C}^n$. Thus, using $1 \in \gcd(f_0(z), \dots, f_m(z))$ we can conclude that there is a character $\chi: H \rightarrow \mathbb{C}^*$ satisfying $f_i(h \cdot x) = \chi(h)f_i(x)$ for every i and every (h, x) in $H \times X$.

Now every f_i is a sum of monomials $q_{i1}, \dots, q_{ir_i}$. Note that also each of the monomials q_{ij} is homogeneous with respect to the character χ . Moreover, since the f_i have no common zero in X , neither have the q_{ij} . Set $r := \sum r_i$ and define a toric morphism

$$g: X \rightarrow \mathbb{P}_r, \quad x \mapsto [q_{01}(x), \dots, q_{0r_0}(x), \dots, q_{m1}(x), \dots, q_{mr_m}(x)].$$

Then g is H -invariant. In order to define an open subset W of $\mathbb{P}_r$ and a regular map $h: W \rightarrow \mathbb{P}_m$ with the desired properties, consider the linear forms

$$L_i: [z_{01}, \dots, z_{0r_0}, \dots, z_{m1}, \dots, z_{mr_m}] \mapsto z_{i1} + \dots + z_{ir_i}$$

on $\mathbb{P}_r$. Set $W := \mathbb{P}_r \setminus V(\mathbb{P}_r; L_1, \dots, L_m)$ and

$$h: W \rightarrow \mathbb{P}_m, \quad [z] \mapsto [L_0(z), \dots, L_m(z)].$$

Since the f_i have no common zero in X we obtain $g(X) \subset W$. Moreover, by construction we have $f = h \circ g$. So the assertion is proved for the case that $Z = \mathbb{P}_m$ and X is an open toric subvariety of some $\mathbb{C}^n$.

In a second step assume that Z is arbitrary but X again is an open toric subvariety of some $\mathbb{C}^n$. Choose a locally closed embedding $\iota: Z \rightarrow \mathbb{P}_m$. By Step one we obtain a decomposition of $f' := \iota \circ f$ as $f' = h' \circ g$, where $g: X \rightarrow \mathbb{P}_r$ is an H -invariant toric morphism such that $g(X)$ is contained in an open subset W' of $\mathbb{P}_r$ and $h': W' \rightarrow \mathbb{P}_m$ is regular.

Then $W := h'^{-1}(\iota(Z))$ is a locally closed subvariety of W' . Moreover we have $g(X) \subset W$ and there is a unique regular map $h: W \rightarrow Z$ with $h' = \iota \circ h$. It follows that $f = h \circ g$ is the desired decomposition.

Finally, let also X be arbitrary. As described in [3], there is an open toric subvariety U of some $\mathbb{C}^n$ and a surjective toric morphism $p: U \rightarrow X$ such that p is the good quotient of U by some algebraic subgroup H_0 of $(\mathbb{C}^*)^n$. Consider $f' := f \circ p$. Then f' is invariant by the action of $H' := \pi^{-1}(H)$, where π denotes the homomorphism of the acting tori associated to p .

By the first two steps we can decompose f' as $f' = h \circ g'$ with an H' -invariant toric morphism $g': U \rightarrow \mathbb{P}_r$ and a regular map $h: W \rightarrow Z$, where $W \subset \mathbb{P}_r$ is locally closed with $g'(U) \subset W$. Since g' is H' -invariant, it is also invariant by the action of H_0 . Thus there

is a unique toric morphism $g: X \rightarrow \mathbb{P}_r$ such that $g' = g \circ p$. Since p is surjective, g is H -invariant and we have $f = h \circ g$ which is the desired decomposition of f . $\square$

Proof of Theorem 1. Let us first assume that the toric quasi-projective reduction $q: X \rightarrow X^{\text{tp}}$ is surjective. Let $f: X \rightarrow Z$ be a regular map to a quasi-projective variety Z . We have to show that f factors uniquely through q .

By Proposition 2.1 there is a toric morphism $g: X \rightarrow X'$ to a projective toric variety X' , and a rational map $h: X' \rightarrow Z$ which is regular on $g(X)$ such that $f = h \circ g$. Now there is a toric morphism $\tilde{g}: X^{\text{tp}} \rightarrow X'$ such that $g = \tilde{g} \circ q$. Since q was assumed to be surjective, we have $\tilde{g}(X^{\text{tp}}) = g(X)$, and hence f factors through q .

Now suppose that $p: X \rightarrow X^{\text{qp}}$ is a quasi-projective reduction. Then clearly p is surjective and X^{qp} is normal. Moreover, there is an induced action of the torus T on X^{qp} making p equivariant. We claim that this action is regular:

According to Proposition 2.1 choose a toric morphism $g: X \rightarrow X'$ to a projective toric variety X' , and a rational map $h: X' \rightarrow X^{\text{qp}}$ such that $g(X)$ is contained in the domain W' of definition of h and $p = h \circ g$. By the universal property of the toric quasi-projective reduction $q: X \rightarrow X^{\text{tp}}$ there is a toric morphism $\tilde{g}: X^{\text{tp}} \rightarrow X'$ such that $g = \tilde{g} \circ q$.

Moreover, by the universal property of p , there is a regular map $\alpha: X^{\text{qp}} \rightarrow X^{\text{tp}}$ such that $q = \alpha \circ p$. Note that $\alpha(X^{\text{qp}}) \subset q(X)$ and $\tilde{g}(q(X)) \subset W'$. So, using surjectivity of p and equivariance of p and q , we obtain for a given pair $(t, y) \in T \times X^{\text{qp}}$ the equality

$$t \cdot y = h(\tilde{g}(t \cdot \alpha(y))).$$

This implies regularity of the induced T -action on X^{qp} . It follows that X^{qp} is in fact a toric variety and p is a toric morphism. Thus we obtain a toric morphism $\beta: X^{\text{tp}} \rightarrow X^{\text{qp}}$ with $p = \beta \circ q$. By uniqueness of the factorizations we obtain that α and β are inverse to each other, i.e., q is also a quasi-projective reduction of X . $\square$

Proof of Theorem 2. Suppose first that $s: X \rightarrow Y$ is a quotient for the action of H on X in the category of quasi-projective varieties. As above we see that Y is a toric variety and s is a surjective toric morphism. The universal property of the toric quotient $s_1: X \rightarrow X_{\text{tor}}^{\text{tor}} H$ yields a toric morphism $q: X_{\text{tor}}^{\text{tor}} H \rightarrow Y$ such that $s = q \circ s_1$. Clearly q satisfies the universal property of the toric quasi-projective reduction of $X_{\text{tor}}^{\text{tor}} H$.

Now let $s_1: X \rightarrow X_{\text{tor}}^{\text{tor}} H$ denote the toric quotient, $q: X_{\text{tor}}^{\text{tor}} H \rightarrow Y$ the toric quasi-projective reduction and assume that $q \circ s_1$ is surjective. Let $f: X \rightarrow Z$ be an H -invariant

regular map to a quasi-projective variety. Choose a decomposition $f = h \circ g$ as in Proposition 2.1. Then, by the universal properties of toric quotient and quasi-projective reduction there is a regular map $\tilde{g}$ with $g = \tilde{g} \circ q \circ s_1$. Since $q \circ s_1$ is surjective, h is defined on $\tilde{g}(Y)$. Thus $f = (h \circ \tilde{g}) \circ (q \circ s_1)$ is the desired factorization of f . $\square$

The above proofs yield in fact the following generalization of Theorems 1 and 2: Let X be any toric variety and let H be an algebraic subgroup of the acting torus T of X . Call an H -invariant regular map $p: X \rightarrow X_H^{\text{qp}}$ to a quasi-projective variety X_H^{qp} an *H -invariant quasi-projective reduction* if it is universal with respect to H -invariant regular maps from X to quasi-projective varieties.

Now write $H = \Gamma H^0$ with a finite subgroup Γ and a subtorus H^0 of T . Let $g: X \rightarrow X'$ denote the geometric quotient for the action of Γ on X . Then g is a toric morphism. Hence there is an induced action of H^0 on X' . Let $s_1: X' \rightarrow X' /_{\text{tor}} H^0$ be the toric quotient for this action and let $q: X' /_{\text{tor}} H^0 \rightarrow Y$ be the quasi-projective toric reduction. Then we obtain:

Theorem 3. *X has an H -invariant quasi-projective reduction if and only $q \circ s_1$ is surjective. If so, then $q \circ s_1 \circ g$ is the H -invariant quasi-projective reduction of X . $\square$*

3 Examples


We first give an example of a 3-dimensional toric variety X_Δ that admits no quasi-projective reduction. This variety is an open toric subvariety of the minimal example for a smooth complete but non-projective toric variety presented in [9], Section 2.3.

3.1 Example. Let e_1, e_2 and e_3 denote the canonical basis vectors of the lattice $\mathbb{Z}^3$. Consider the vectors

$$\begin{aligned} v_1 &:= -e_1, & v'_1 &:= e_2 + e_3, \\ v_2 &:= -e_2, & v'_2 &:= e_1 + e_3, \\ v_3 &:= -e_3, & v'_3 &:= e_1 + e_2. \end{aligned}$$


Let Δ be the fan in $\mathbb{Z}^3$ with the maximal cones

$$\tau_1 := \text{cone}(v_1, v'_3), \quad \tau_2 := \text{cone}(v_2, v'_1) \quad \text{and} \quad \tau_3 := \text{cone}(v_3, v'_2).$$



We claim that the toric quasi-projective reduction q of X_Δ is the toric morphism associated to id_N interpreted as a map from Δ to the fan $\tilde{\Delta}$ having as its maximal cones

$$\sigma_1 := \text{cone}(v_1, v_3, v'_1, v'_3), \quad \sigma_2 := \text{cone}(v_1, v_2, v'_1, v'_2) \quad \text{and} \quad \sigma_3 := \text{cone}(v_2, v_3, v'_2, v'_3).$$



Note that q is not surjective. In order to prove that q is the toric quasi-projective reduction of X_Δ , we have to show that every Δ -concave family $(u_i)_{i=1,2,3}$ can be extended to a $\tilde{\Delta}$ -concave family. Note that $v_1 + v'_3$ equals $v_3 + v'_1$ and hence we have

$$u_1(v_1) + u_1(v'_3) = u_1(v_3) + u_1(v'_1) \geq u_3(v_3) + u_2(v'_1).$$

Similarly we obtain

$$\begin{aligned} u_2(v_2) + u_2(v'_1) &= u_2(v_1) + u_2(v'_2) \geq u_1(v_1) + u_3(v'_2), \\ u_3(v_3) + u_3(v'_2) &= u_3(v'_3) + u_3(v_2) \geq u_1(v'_3) + u_2(v_2). \end{aligned}$$

Summing over these three inequalities, we arrive at an identity, and therefore the inequalities are in fact equalities. This implies

$$\begin{aligned} u_1(v_1) &= u_2(v_1), & u_1(v'_1) &= u_2(v'_1), \\ u_1(v_3) &= u_3(v_3), & u_1(v'_3) &= u_3(v'_3), \\ u_2(v_2) &= u_3(v_2), & u_2(v'_2) &= u_3(v'_2). \quad \diamond \end{aligned}$$

In the above example the quasi-projective toric reduction has a trivial kernel, and the variety $X_{\tilde{\Delta}}$ has the same dimension as X_Δ . For the complete case we have more generally:

3.2 Remark. Let Δ be a complete fan in a lattice N . Then $\dim X_\Delta = \dim X_\Delta^{\text{qp}}$ holds if and only if Δ can be defined via a subdivision of a lattice polytope in $N_{\mathbb{R}}$. $\square$

The next example is taken from the book of Fulton. It shows that in general a complete toric variety is very far from its projective reduction.

3.3 Example. Consider the complete fan in $\mathbb{Z}^3$ obtained by taking the cones over the faces of the standard cube with vertices $(\pm 1, \pm 1, \pm 1)$. Deform this fan into a new complete fan Δ in $\mathbb{Z}^3$ by moving the vertex $(1, 1, 1)$ to $(1, 2, 3)$. The only support functions of Δ are the linear functions in M (see [6], p. 26). So X_Δ^{qp} is a point. $\diamond$


Now we turn to quotients of a quasi-projective toric variety X with acting torus T by subtori $H \subset T$. Examples of such quotients are obtained by Mumford's Geometric Invariant Theory:

For the sake of simplicity assume $X = \mathbb{P}_n$. Then the choice of a lifting of the T -action to $\mathbb{C}^{n+1}$ yields a notion of H -semistability. The set $X^{\text{ss}} \subset X$ of H -semistable points is T -invariant and there is a quotient $X^{\text{ss}} \rightarrow Y$ in the category of quasi-projective varieties for the action of H on X^{ss} (see [8], also [7] and [2]).

3.4 Example. Let Δ be the fan in $\mathbb{R}^4$ that has $\sigma_1 := \text{cone}(e_1, e_2)$ and $\sigma_2 := \text{cone}(e_3, e_4)$ as maximal cones. Then X_Δ is an open toric subvariety of $\mathbb{C}^4$ with acting torus $T = \mathbb{C}^{*4}$. Define a projection $S_1: \mathbb{Z}^4 \rightarrow \mathbb{Z}^3$ by setting

$$S_1(e_1) := e_1, \quad S_1(e_2) := e_2, \quad S_1(e_3) := e_3, \quad S_1(e_4) := e_1 + e_2.$$

Then $S_1(e_1), \dots, S_1(e_4)$ generate $\tau := \text{cone}(e_1, e_2, e_3) \subset \mathbb{R}^3$. The faces $\text{cone}(e_1, e_3)$ and $\text{cone}(e_2, e_3)$ of τ are not contained in $S_1(|\Delta|)$.



By [1], the toric morphism $s_1: X_\Delta \rightarrow X_\tau$ defined by S_1 is the toric quotient for the action of the subtorus $H \subset T$ corresponding to the sublattice $\ker(S_1)$ of $\mathbb{Z}^4$. In particular, s_1 is not surjective. So the action of H on X_Δ has no quotient in the category of quasi-projective varieties. $\diamond$

Note that surjectivity of the toric quotient $s_1: X \rightarrow X_{\text{tor}}/H$ does not imply the existence of a quotient in the category of quasi-projective varieties:

3.5 Example. Let Δ' be the fan in $\mathbb{R}^3$ with the maximal cones

$$\tau_1 := \text{cone}(e_1, e_2), \quad \tau_2 := \text{cone}(e_3, e_4), \quad \tau_3 := \text{cone}(e_5, e_6).$$

Then the associated toric variety $X_{\Delta'}$ is an open toric subvariety of $\mathbb{C}^6$. In the notation of Example 3.1, define a projection $S_1: \mathbb{Z}^6 \rightarrow \mathbb{Z}^3$ by

$$\begin{aligned} S_1(e_1) &:= -e_1, & S_1(e_2) &:= v'_1, \\ S_1(e_3) &:= -e_2, & S_1(e_4) &:= v'_2, \\ S_1(e_5) &:= -e_3, & S_1(e_6) &:= v'_3. \end{aligned}$$

Then S_1 is a map of the fan Δ' and the fan Δ of 3.1, in fact the (surjective) toric morphism $s_1: X_{\Delta'} \rightarrow X_{\Delta}$ associated to S_1 is the toric quotient of the action of the subtorus H of $\mathbb{C}^{*6}$ corresponding to the sublattice $\ker(S_1) \subset \mathbb{Z}^6$. Since the quasi-projective reduction of X_{Δ} is not surjective, there is no quotient in the category of quasi-projective varieties for the action of H on $X_{\Delta'}$.

References

- [1] A. A'Campo-Neuen, J. Hausen, Quotients of Toric Varieties by the Action of a Subtorus, to appear in Tôhoku Math. J..
- [2] A. Białynicki-Birula, J. Świąćicka, Open Subsets of Projective Spaces with a Good Quotient by an Action of a Reductive Group, Transformation Groups Vol.1, No. 3 (1996), 153-185.
- [3] D. Cox, The Homogeneous Coordinate Ring of a Toric Variety, J. Algebraic Geometry **4** (1995), 17-51.
- [4] A. Grothendieck, J. Dieudonné, Éléments de géométrie algébrique, Chap. II, Étude globale élémentaire de quelques classes de morphismes, Publications mathématiques de l'IHES, n. 8, 1961.
- [5] G. Ewald, Combinatorial Convexity and Algebraic Geometry, Springer, New York, 1996.

- [6] *W. Fulton*, Introduction to Toric Varieties, Princeton University Press, Princeton, 1993.
- [7] *M. Kapranov, B. Sturmfels, A. V. Zelevinsky*, Quotients of Toric Varieties, Math. Ann. **290** (1991), 643-655.
- [8] *D. Mumford, J. Fogarty, F. Kirwan*, Geometric Invariant Theory, 3rd enlarged edition, Springer, Berlin, 1994.
- [9] *T. Oda*, Convex Bodies and Algebraic Geometry, Springer, Berlin, 1988.

FAKULTÄT FÜR MATHEMATIK UND INFORMATIK
UNIVERSITÄT KONSTANZ
FACH D197
D-78457 KONSTANZ
GERMANY

E-mail address: `Annette.ACampo@uni-konstanz.de`
`Juergen.Hausen@uni-konstanz.de`